Example text for anonymous proposing

There is some apprehension on how to write the body of the PDF attachment. The following examples keep to the Proposer Guidelines in Anonymous Reviews, and are meant as additional guides in crafting your own proposals.

Here is an example of text from a sample proposal:

Over the last five years, we have used infrared photometry from 2MASS to compile a census of nearby ultracool M and L dwarfs (Cruz et al, 2003; 2006). We have identified 87 L dwarfs in 80 systems with nominal distances less than 20 parsecs from the Sun. This is the first true L dwarf census – a large-scale, volume-limited sample. Most distances are based on spectroscopic parallaxes, accurate to 20%, which is adequate for present purposes. Fifty systems already have high-resolution imaging, including our Cycle 9 and 13 snapshot programs, #8581 and #10143; nine are in binary or multiple systems, including six new discoveries. We propose to target the remaining sources via the current proposal. We will combine those data with X-ray measurements made in our Cycle 14 Chandra proposal #2095.

Here is the same text, re-worked following the anonymizing guidelines:

Over the last five years, 2MASS infrared photometry has been used to compile a census of nearby ultracool M and L dwarfs (Cruz et al, 2003; 2006). 87 L dwarfs in 80 systems have been identified with nominal distances less than 20 parsecs from the Sun. This is the first true L dwarf census — a large-scale, volume-limited sample. Most distances are based on spectroscopic parallaxes, accurate to 20%, which is adequate for present purposes. Fifty systems already have high-resolution imaging, including the Cycle 9 and 13 snapshot programs, #8581 and #10143; nine are in binary or multiple systems, including six new discoveries. We propose to target the remaining sources via the current proposal. Those data will be combined with Chandra X-ray measurements from program #2095 (Reid, priv. comm.).

Here is another example of text from a sample proposal:

In Rogers et al. (2014), we concluded that the best explanation for the dynamics of the shockwave and the spectra from both the forward-shocked ISM and the reverse-shocked ejecta is that a Type Ia supernova exploded into a preexisting wind-blown cavity. This object is the only known example of such a phenomenon, and it thus provides a unique opportunity to illuminate the nature of Type Ia supernovae and the progenitors. If our model from Rogers et al. (2014) is correct, then the single-degenerate channel for SNe Ia production must exist. We propose here for a second epoch of observations which we will compare with our first epoch obtained in 2007 to measure the proper motion of the shock wave.

Here is the same text, again re-worked following the anonymizing guidelines:

Rogers et al. (2014) concluded that the best explanation for the dynamics of the shockwave and the spectra from both the forward-shocked ISM and the reverse-shocked ejecta is that a Type Ia supernova exploded into a preexisting wind-blown cavity. This object is the only known example of such a phenomenon, and it thus provides a unique opportunity to illuminate the nature of Type Ia supernovae and the progenitors. If the model from Rogers et al. (2014) is correct, then the single-degenerate channel for SNe Ia production must exist. We propose here for a second epoch of observations which we will compare with a first epoch obtained in 2007 to measure the proper motion of the shock wave.

As further example, here's an excerpted sample from an Archival proposal:

